


We claim:


In the following claims the subscript and superscripts of a given variable are distinct.
For example, R₁ is distinct from R¹.

5 1. A non-nucleoside reverse transcriptase inhibitor compound comprising a
phosphonate group.





2. A non-nucleoside reverse transcriptase inhibitor of claim 1 selected from:
Capravirine-like phosphonate NNRTI compounds,
10 PETT-like phosphonate NNRTI compounds,
Pyrazole-like phosphonate NNRTI compounds,
Urea-PETT-like phosphonate NNRTI compounds,
Nevirapine-like phosphonate NNRTI compounds,
Quinazolinone-like phosphonate NNRTI compounds,
15 Efavirenz-like phosphonate NNRTI compounds,
Benzophenone-like phosphonate NNRTI compounds,
Pyrimidine-like phosphonate NNRTI compounds,
SJ3366-like phosphonate NNRTI compounds,
Delavirdine-like phosphonate NNRTI compounds,
20 Emivirine-like phosphonate NNRTI compounds,
Loviride-like phosphonate NNRTI compounds, and
UC781-like phosphonate NNRTI compounds;
and pharmaceutically acceptable salts, hydrates, and formulations thereof.

25 3. A compound according to claim 1 selected from the Formulas:







where Z is CH or N,



5


10




where A is O, S, or N(R^x) and X is O or S;

5 wherein A⁰ is A¹, A² or W³ with the proviso that the compound includes at least one A¹;


A¹ is:



A² is:



A³ is:




Y^1 is independently O, S, $N(R^x)$, $N(O)(R^x)$, $N(OR^x)$, $N(O)(OR^x)$, or $N(N(R^x)(R^x))$;

Y^2 is independently a bond, O, $N(R^x)$, $N(O)(R^x)$, $N(OR^x)$, $N(O)(OR^x)$, $N(N(R^x)(R^x))$,

$-S(O)_{M2}-$, or $-S(O)_{M2}-S(O)_{M2}-$;

5 R^x is independently H, R^1 , W^3 , a protecting group, or the formula:



R^y is independently H, W^3 , R^2 or a protecting group;

R^1 is independently H or alkyl of 1 to 18 carbon atoms;

R^2 is independently H, R^1 , R^3 or R^4 wherein each R^4 is independently substituted with

10 0 to 3 R^3 groups, or taken together at a carbon atom, two R^2 groups form a ring of 3 to 8 carbons and the ring may be substituted with 0 to 3 R^3 groups;

R^3 is R^{3a} , R^{3b} , R^{3c} or R^{3d} , provided that when R^3 is bound to a heteroatom, then R^3 is R^{3c} or R^{3d} ;

R^{3a} is F, Cl, Br, I, $-CN$, N_3 or $-NO_2$;

15 R^{3b} is Y^1 ;

R^{3c} is $-R^x$, $-N(R^x)(R^x)$, $-SR^x$, $-S(O)R^x$, $-S(O)_2R^x$, $-S(O)(OR^x)$, $-S(O)_2(OR^x)$, $-OC(Y^1)R^x$, $-OC(Y^1)OR^x$, $-OC(Y^1)(N(R^x)(R^x))$, $-SC(Y^1)R^x$, $-SC(Y^1)OR^x$, $-SC(Y^1)(N(R^x)(R^x))$, $-N(R^x)C(Y^1)R^x$, $-N(R^x)C(Y^1)OR^x$, or $-N(R^x)C(Y^1)(N(R^x)(R^x))$;

R^{3d} is $-C(Y^1)R^x$, $-C(Y^1)OR^x$ or $-C(Y^1)(N(R^x)(R^x))$;

20 R^4 is an alkyl of 1 to 18 carbon atoms, alkenyl of 2 to 18 carbon atoms, or alkynyl of 2 to 18 carbon atoms;

R^5 is R^4 wherein each R^4 is substituted with 0 to 3 R^3 groups;

W^3 is W^4 or W^5 ;

W^4 is R^5 , $-C(Y^1)R^5$, $-C(Y^1)W^5$, $-SO_2R^5$, or $-SO_2W^5$;

W^5 is carbocycle or heterocycle wherein W^5 is independently substituted with 0 to 3 R^2

groups;

W^6 is W^3 are independently substituted with 1, 2, or 3 A^3 groups;

$M2$ is 0, 1 or 2;


$M12a$ is 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12;

$M12b$ is 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12.


$M1a$, $M1c$, and $M1d$ are independently 0 or 1; and

$M12c$ is 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12.


4. A compound of claim 3 having Formula Ia:




5. A compound of claim 3 having Formula Ib:




6. A compound of claim 3 having Formula Ic:



7. A compound of claim 3 having Formula Id:




8. A compound of claim 3 having the formulas:




5


9. A compound of claim 3 having the formulas:



10. A compound of claim 3 having the formulas:




11. A compound of claim 3 having the formulas:




5

12. A compound of claim 3 having the formulas:




13. A compound of claim 3 having the formulas:




5


14. A compound of claim 3 having the formulas:




15. A compound of claim 3 having the formulas:




16. A compound of claim 3 having the formulas:



17. A compound of claim 3 having the formulas:




18. A compound of claim 3 having the formulas:




5

19. A compound of claim 3 having the formulas:




20. The compound of claim 3 wherein A^1 is of the formula:




5

21. The compound of claim 20 wherein A^1 is of the formula:




22. The compound of claim 21 wherein A^1 is of the formula:



5

23. The compound of claim 22 wherein A^1 is of the formula:








and W^{5a} is a carbocycle or a heterocycle where W^{5a} is independently substituted with 0 or 1 R^2 groups.

10


24. The compound of claim 23 wherein M12a is 1.

25. The compound of claim 23 wherein A^1 is selected from the formulas:




R_1 and R_2 are independently selected from hydroxy, methoxy, ethoxy, trifluoroethoxy, isopropoxy, phenoxy, benzyloxy, O-pivaloyloxymethyl, an amino acid ester and a lactate ester; and

5 W^{5a} is selected from the formulas:




26. The compound of claim 3 wherein A³ is of the formula:




5

27. The compound of claim 26 wherein A³ is of the formula:




28. The compound of claim 27 wherein A^3 is of the formula:



Y^{1a} is O or S; and


Y^{2a} is O, N(R^x) or S.

29. The compound of claim 28 wherein A^3 is of the formula:



and Y^{2b} is O or N(R^x).


30. The compound of claim 29 wherein A^3 is of the formula:



Y^{2b} is O or N(R^x); and

M12d is 1, 2, 3, 4, 5, 6, 7 or 8.

31. The compound of claim 30 wherein A^3 is of the formula:




Y^{2b} is O or $N(R^x)$; and


M12d is 1, 2, 3, 4, 5, 6, 7 or 8.

32. The compound of claim 31 wherein M12d is 1.

33. The compound of claim 3 wherein A^3 is of the formula:




34. The compound of claim 33 wherein A^3 is of the formula:



35. The compound of claim 34 wherein W^5 is a carbocycle.

5

36. The compound of claim 35 wherein A^3 is of the formula:




37. The compound of claim 36 wherein W^5 is phenyl.

10

38. The compound of claim 33 wherein M12b is 1.


39. The compound of claim 38 wherein A^3 is of the formula:



Y^{1a} is O or S; and


Y^{2a} is O, N(R^x) or S.

40. The compound of claim 39 wherein A³ is of the formula:



and Y^{2b} is O or N(R^x).

41. The compound of claim 40 wherein A³ is of the formula:




Y^{2b} is O or N(R^x); and

M12d is 1, 2, 3, 4, 5, 6, 7 or 8.

42. The compound of claim 41 wherein R¹ is H.

43. The compound of claim 41 wherein M12d is 1.


44. The compound of claim 41 wherein A³ is of the formula:




wherein the phenyl carbocycle is substituted with 0 to 3 R^2 groups, and R^1 is H or C_1-C_{18} alkyl.

5 45. The compound of claim 44 wherein Y^{2b} is $N(R^x)$.


46. The compound of claim 45 wherein A^3 is of the formula:




10 47. The compound of claim 44 wherein A^3 is of the formula:



48. A compound of claim 3 wherein R^x is of the formula:



49. A compound of claim 48 wherein R^x is of the formula:




5

Y^{1a} is O or S; and

Y^{2c} is O, N(R^y) or S.


50. A compound of claim 48 wherein R^x is of the formula:




10

and Y^{2d} is O or N(R^y).


51. A compound of claim 50 wherein R^x is of the formula:



52. A compound of claim 51 wherein R^x is of the formula:




53. The compound of claim 3 wherein R^x is of the formula:




5

54. The compound of claim 53 wherein A^3 is of the formula:




10

55. The compound of claim 3 wherein A^3 is of the formula:




; and

R^x is of the formula:



56. The compound of claim 55 wherein A³ is of the formula:




5

Y^{1a} is O or S; and

Y^{2a} is O, N(R²) or S.

57. The compound of claim 56 wherein A³ is of the formula:




10

Y^{2b} is O or N(R²); and

Y^{2c} is O, N(R^y) or S.

15

58. The compound of claim 57 wherein A³ is of the formula:




Y^{1a} is O or S;

Y^{2b} is O or $N(R^2)$;

Y^{2d} is O or $N(R^y)$; and

M12d is 1, 2, 3, 4, 5, 6, 7 or 8.


59. The compound of claim 58 wherein A^3 is of the formula:



Y^{2b} is O or $N(R^2)$; and


M12d is 1, 2, 3, 4, 5, 6, 7 or 8.

60. The compound of claim 59 wherein A^3 is of the formula:




and Y^{2b} is O or $N(R^2)$.


61. The compound of claim 60 wherein A^3 is of the formula:




5 62. The compound of claim 3 wherein A^3 is of the formula:



R^x is of the formula:




10 63. The compound of claim 62 wherein A^3 is of the formula:



Y^{1a} is O or S; and

Y^{2a} is O, N(R²) or S.


64. The compound of claim 63 wherein A³ is of the formula:



Y^{2b} is O or N(R²); and

Y^{2c} is O, N(R^y) or S.

65. The compound of claim 64 wherein A³ is of the formula:



R¹ is H or C₁–C₁₈ alkyl;


Y^{2d} is O or N(R^y); and

M12d is 1, 2, 3, 4, 5, 6, 7 or 8.


66. The compound of claim 65 wherein R¹ is H; and Y^{1a} and Y^{2d} are O.

67. The compound of claim 66 wherein M12d is 1.

68. The compound of claim 3 wherein A¹ is of the formula:




A³ is of the formula:




5


69. The compound of claim 68 wherein A¹ is of the formula:



A³ is of the formula:




R^x is of the formula:




5

70. The compound of claim 69 wherein A^1 is of the formula:



A^3 is of the formula:




10

Y^{1a} is O or S; and


Y^{2a} is O, N(R²) or S.

5 71. The compound of claim 70 wherein A¹ is of the formula:



W^{5a} is a carbocycle independently substituted with 0 or 1 R² groups;

A³ is of the formula:




10 Y^{1a} is O or S;

Y^{2b} is O or N(R²); and


Y^{2c} is O, N(R^y) or S.

72. The compound of claim 71 wherein A¹ is of the formula:



W^{5a} is a carbocycle independently substituted with 0 or 1 R² groups;

A³ is of the formula:




Y^{1a} is O or S;

Y^{2b} is O or $N(R^2)$;

Y^{2d} is O or $N(R^y)$; and

M12d is 1, 2, 3, 4, 5, 6, 7 or 8.


73. The compound of claim 72 wherein A^1 is of the formula:



Y^{2b} is O or $N(R^2)$; and


M12d is 1, 2, 3, 4, 5, 6, 7 or 8.

74. The compound of claim 3 wherein A^1 is of the formula:




; and

A³ is of the formula:




75. The compound of claim 74 wherein A¹ is of the formula:




5

A³ is of the formula:




10

R^x is of the formula:




76. The compound of claim 75 wherein A¹ is of the formula:



5

A³ is of the formula:




Y^{1a} is O or S; and

Y^{2a} is O, N(R²) or S.


10

77. The compound of claim 76 wherein A¹ is of the formula:



W^{5a} is a carbocycle independently substituted with 0 or 1 R^2 groups;

A^3 is of the formula:




5

Y^{1a} is O or S;

Y^{2b} is O or $N(R^2)$; and

Y^{2c} is O, $N(R^y)$ or S.


78. The compound of claim 77 wherein A^3 is of the formula:



10


wherein the phenyl carbocycle is substituted with 0 to 3 R^2 groups.

79. The compound of claim 75 wherein A^1 is of the formula:



W^{5a} is a carbocycle or heterocycle where W^{5a} is independently substituted with 0 or 1 R^2 groups;

5 A^3 is of the formula:




Y^{1a} is O or S;

Y^{2b} is O or $N(R^2)$;


Y^{2d} is O or $N(R^y)$; and

10 M12d is 1, 2, 3, 4, 5, 6, 7 or 8.


80. The compound of claim 79 wherein A^1 is of the formula:



15 81. The compound of claim 3 wherein A^2 is of the formula:



82. The compound of claim 81 wherein A^2 is of the formula:




5

83. The compound of claim 82 wherein M12b is 1.

84. The compound of claim 82 where M12b is 0, Y^2 is a bond and W^5 is a carbocycle or heterocycle where W^5 is optionally and independently substituted with 1, 2, or 3 R^2 groups.

10

85. The compound of claim 82 wherein A^2 is of the formula:



and W^{5a} is a carbocycle or heterocycle where W^{5a} is optionally and independently substituted with 1, 2, or 3 R^2 groups.


15

86. The compound of claim 85 wherein M12a is 1.


87. The compound of claim 86 wherein A² is selected from phenyl, substituted phenyl, benzyl, substituted benzyl, pyridyl and substituted pyridyl.

5

88. The compound of claim 3 wherein A² is of the formula:




89. The compound of claim 88 wherein A² is of the formula:



10


90. The compound of claim 89 wherein M12b is 1.

91. The compound of claim 6 having the formula:




15

wherein A¹ is of the formula:



; and

A³ is of the formula:




5

92. The compound of claim 91 wherein A² is selected from phenyl, substituted phenyl, benzyl, substituted benzyl, pyridyl and substituted pyridyl.


93. The compound of claim 92 wherein substituted phenyl, substituted benzyl, and substituted pyridyl are independently substituted with 1, 2 or 3 R² groups.

10

94. The compound of claim 91 wherein A¹ is of the formula:




A³ is of the formula:




5

95. The compound of claim 94 wherein A³ is of the formula:




96. The compound of claim 94 wherein R^x is of the formula:




10


97. The compound of claim 96 wherein A³ is of the formula:




98. The compound of claim 95 wherein R^x is of the formula:



99. The compound of claim 98 wherein A³ is of the formula:




100. The compound of claim 91 having the formula:




101. The compound of claim 100 wherein R⁴ is isopropyl.

102. The compound of claim 100 having the formula:




103. The compound of claim 100 wherein A¹ is of the formula:



5


A³ is of the formula:



and Y^{1a} is O or S.


10

104. The compound of claim 103 wherein A³ is of the formula:



and Y^{2a} is O, N(R^2) or S.


105. The compound of claim 104 wherein A^3 is of the formula:



Y^{2b} is O or N(R^2); and

Y^{2c} is O, N(R^y) or S.

106. The compound of claim 105 wherein A^3 is of the formula:




Y^{1a} is O or S;

Y^{2b} is O or N(R^2);

Y^{2d} is O or N(R^y); and


M12d is 1, 2, 3, 4, 5, 6, 7 or 8.

107. The compound of claim 106 wherein A¹ is of the formula:




5 Y^{2b} is O or N(R²); and
M12d is 1, 2, 3, 4, 5, 6, 7 or 8.

108. The compound of claim 106 wherein A¹ is of the formula:




10 and Y^{2b} is O or N(R²); and
M12d is 1, 2, 3, 4, 5, 6, 7 or 8.


109. The compound of claim 102 wherein W⁵ is selected from the structures:



110. The compound of claim 91 wherein A^1 is of the formula:



5 n is an integer from 1 to 18; A^3 is of the formula:




and Y^{2c} is O, $N(R^y)$ or S.


111. The compound of claim 110 wherein R^1 is H and n is 1.

5

112. The compound of claim 5 having the formula:




wherein A^1 is of the formula:




10


and A^3 is of the formula:




113. The compound of claim 112 wherein A¹ is of the formula:




5 A³ is of the formula:



114. The compound of claim 113 wherein A³ is of the formula:




115. The compound of claim 113 wherein R^x is of the formula:




5

116. The compound of claim 115 wherein A^3 is of the formula:




117. The compound of claim 113 wherein R^x is of the formula:




10

118. The compound of claim 117 wherein A^3 is of the formula:




119. The compound of claim 112 wherein A^2 is selected from:




5 where W^5 is a carbocycle or a heterocycle and where W^5 is independently substituted with 0 to 3 R^2 groups.

120. The compound of claim 112 having the formula:




10

wherein A^1 is of the formula:




A^3 is of the formula:




and Y^{1a} is O or S.

121. The compound of claim 120 wherein A¹ is of the formula:



5


122. The compound of claim 120 wherein A³ is of the formula:



and Y^{2a} is O, N(R²) or S.


10

123. The compound of claim 112 wherein A³ is of the formula:




and Y^{2c} is O, $N(R^y)$ or S.

124. The compound of claim 112 wherein A^1 is of the formula:



5

A^3 is of the formula:




W^{5a} is a carbocycle or a heterocycle where the carbocycle or heterocycle is

10 independently substituted with 0 to 3 R^2 groups;


Y^{2b} is O or $N(R^2)$; and

Y^{2c} is O, $N(R^y)$ or S.

125. The compound of claim 124 wherein A^1 is of the formula:



A³ is of the formula:



5

Y^{1a} is O or S;


Y^{2b} is O or N(R²);

Y^{2d} is O or N(R^y); and

M12d is 1, 2, 3, 4, 5, 6, 7 or 8.

10

126. The compound of claim 125 wherein A¹ is of the formula:




15

Y^{2b} is O or N(R²); and

M12d is 1, 2, 3, 4, 5, 6, 7 or 8.

127. The compound of claim 126 wherein A^1 is of the formula:




and Y^{2b} is O or $N(R^2)$; and

M12d is 1, 2, 3, 4, 5, 6, 7 or 8.


128. The compound of claim 112 wherein A^2 is a phenyl substituted with 0 to 3 R^2 groups.

129. The compound of claim 120 wherein W^4 is of the formula:




wherein n is an integer from 1 to 18; and Y^{2b} is O or $N(R^2)$.


130. The compound of claim 129 having the formula:




131. A compound of claim 130 selected from the formulas:









132. A compound of claim 6 selected from the structures:



17



18






5


10

and


133. A compound of claim 4 selected from the structures:






and





134. A compound of claim 6 selected from the structures:








5

135. A compound of claim 5 selected from the structures:




10








5






and



136. A compound of claim 5 selected from the structures:

10






5


10

and




78.


137. A compound of claim 5 selected from the structures:




81




82




83




84




85




86



87




88




89

5

10





and





5

138. A compound of claim 5 selected from the structures:




10






108




110




5



115




117




5


and



139. A compound of claim 3 selected from the structures:




97




97a

and



122 .

140. A compound of the formula:



wherein:

A₁ is -(X₂-(C(R₂)(R₂))_{m1}-X₃)_{m1}-W₃, wherein W₃ is substituted with 1 to 3 A₃ groups;

A₂ is -(X₂-(C(R₂)(R₂))_{m1}-X₃)_{m1}-W₃;

A₃ is -(X₂-(C(R₂)(R₂))_{m1}-X₃)_{m1}-P(Y₁)(Y₁R_{6a})(Y₁R_{6a});

X₂ and X₃ are independently a bond, -O-, -N(R₂)-, -N(OR₂)-, -N(N(R₂)(R₂))- , -S-,

-SO-, or -SO₂-;

each Y₁ is independently O, N(R₂), N(OR₂), or N(N(R₂)(R₂)), wherein each Y₁ is bound by two single bonds or one double bond;

R₁ is independently H or alkyl of 1 to 12 carbon atoms;

R₂ is independently H, R₃ or R₄ wherein each R₄ is independently substituted with 0 to 3 R₃ groups;

R₃ is independently F, Cl, Br, I, -CN, N₃, -NO₂, -OR_{6a}, -OR₁, -N(R₁)₂, -N(R₁)(R_{6b}), -N(R_{6b})₂, -SR₁, -SR_{6a}, -S(O)R₁, -S(O)₂R₁, -S(O)OR₁, -S(O)OR_{6a}, -S(O)₂OR₁, -S(O)₂OR_{6a}, -C(O)OR₁, -C(O)R_{6c}, -C(O)OR_{6a}, -OC(O)R₁, -N(R₁)(C(O)R₁), -N(R_{6b})(C(O)R₁), -N(R₁)(C(O)OR₁), -N(R_{6b})(C(O)OR₁), -C(O)N(R₁)₂, -C(O)N(R_{6b})(R₁), -C(O)N(R_{6b})₂, -C(NR₁)(N(R₁)₂), -C(N(R_{6b}))(N(R₁)₂), -C(N(R₁))(N(R₁)(R_{6b})), -C(N(R_{6b}))(N(R₁)(R_{6b})), -C(N(R₁))(N(R_{6b})₂), -C(N(R_{6b}))(N(R_{6b})₂), -N(R₁)C(N(R₁))(N(R₁)₂), -N(R₁)C(N(R₁))(N(R₁)(R_{6b})), -N(R₁)C(N(R_{6b}))(N(R₁)₂), -N(R_{6b})C(N(R₁))(N(R₁)₂), -N(R_{6b})C(N(R_{6b}))(N(R₁)₂), -N(R_{6b})C(N(R₁))(N(R₁)(R_{6b})), -N(R₁)C(N(R₁))(N(R_{6b})₂), -N(R_{6b})C(N(R_{6b}))(N(R₁)(R_{6b})), -N(R_{6b})C(N(R₁))(N(R_{6b})₂), -N(R₁)C(N(R_{6b}))(N(R_{6b})₂), -N(R_{6b})C(N(R_{6b}))(N(R_{6b})₂), =O, =S, =N(R₁), =N(R_{6b}) or W₅;

R₄ is independently alkyl of 1 to 12 carbon atoms, alkenyl of 2 to 12 carbon atoms, or alkynyl of 2 to 12 carbon atoms;

R₅ is independently R₄ wherein each R₄ is substituted with 0 to 3 R₃ groups;

R_{5a} is independently alkylene of 1 to 12 carbon atoms, alkenylene of 2 to 12 carbon atoms, or alkynylene of 2-12 carbon atoms any one of which alkylene, alkenylene or alkynylene is substituted with 0-3 R₃ groups;

R_{6a} is independently H or an ether- or ester-forming group;

R_{6b} is independently H, a protecting group for amino or the residue of a carboxyl-containing compound;

R_{6c} is independently H or the residue of an amino-containing compound;

W₃ is W₄ or W₅;

W₄ is R₅, -C(Y₁)R₅, -C(Y₁)W₅, -SO₂R₅, or -SO₂W₅;

W₅ is carbocycle or heterocycle wherein W₅ is independently substituted with 0 to 3 R₂ groups;

m₁ is independently an integer from 0 to 12, wherein the sum of all m₁'s within each individual claim of A₁, A₂ or A₃ is 12 or less; and

5 m₂ is independently an integer from 0 to 2.


141. The compound of claim 140 wherein:

A₁ is -(C(R₂)(R₂))_{m1}-W₃, wherein W₃ is substituted with 1 A₃ group;


A₂ is -(C(R₂)(R₂))_{m1}-W₃; and

10 A₃ is -(C(R₂)(R₂))_{m1}-P(Y₁)(Y₁R_{6a})(Y₁R_{6a}).


142. The compound of claim 141 having the formula:




15 143. The compound of claim 142 having the formula:



144. The compound of claim 143 having the formula:




145. The compound of claim 144 having the formula:




5


146. The compound of claim 145 wherein W⁵ is selected from:



147. A compound of claim 140 having the formula:



148. A compound of the formula:



5

wherein:

A₁ is -(X₂-(C(R₂)(R₂))_{m1}-X₃)_{m1}-W₃, wherein W₃ is substituted with 1 to 3 A₃

groups;

A₂ is -(X₂-(C(R₂)(R₂))_{m1}-X₃)_{m1}-W₃;

A₃ is -(X₂-(C(R₂)(R₂))_{m1}-X₃)_{m1}-P(Y₁)(Y₁R_{6a})(Y₁R_{6a});

10

X₂ and X₃ are independently a bond, -O-, -N(R₂)-, -N(OR₂)-, -N(N(R₂)(R₂))-, -S-, -SO-, or -SO₂-;

each Y₁ is independently O, N(R₂), N(OR₂), or N(N(R₂)(R₂)), wherein each Y₁ is bound by two single bonds or one double bond;

R₁ is independently H or alkyl of 1 to 12 carbon atoms;

15

R₂ is independently H, R₁, R₃ or R₄ wherein each R₄ is independently substituted with 0 to 3 R₃ groups;

R₃ is independently F, Cl, Br, I, -CN, N₃, -NO₂, -OR_{6a}, -OR₁, -N(R₁)₂, -N(R₁)(R_{6b}), -N(R_{6b})₂, -SR₁, -SR_{6a}, -S(O)R₁, -S(O)₂R₁, -S(O)OR₁, -S(O)OR_{6a}, -S(O)₂OR₁, -S(O)₂OR_{6a}, -C(O)OR₁, -C(O)R_{6c}, -C(O)OR_{6a}, -OC(O)R₁, -N(R₁)(C(O)R₁), -N(R_{6b})(C(O)R₁), -N(R₁)(C(O)OR₁), -N(R_{6b})(C(O)OR₁), -C(O)N(R₁)₂,

20

-C(O)N(R6b)(R1), -C(O)N(R6b)2, -C(NR1)(N(R1)2), -C(N(R6b))(N(R1)2),
 -C(N(R1))(N(R1)(R6b)), -C(N(R6b))(N(R1)(R6b)), -C(N(R1))(N(R6b)2),
 -C(N(R6b))(N(R6b)2), -N(R1)C(N(R1))(N(R1)2), -N(R1)C(N(R1))(N(R1)(R6b)),
 -N(R1)C(N(R6b))(N(R1)2), -N(R6b)C(N(R1))(N(R1)2), -N(R6b)C(N(R6b))(N(R1)2),
 5 -N(R6b)C(N(R1))(N(R1)(R6b)), -N(R1)C(N(R6b))(N(R1)(R6b)),
 -N(R1)C(N(R1))(N(R6b)2), -N(R6b)C(N(R6b))(N(R1)(R6b)), -N(R6b)C(N(R1))(N(R6b)2),
 -N(R1)C(N(R6b))(N(R6b)2), -N(R6b)C(N(R6b))(N(R6b)2), =O, =S, =N(R1), =N(R6b) or
 W5;

R4 is independently alkyl of 1 to 12 carbon atoms, alkenyl of 2 to 12 carbon atoms, or
 10 alkynyl of 2 to 12 carbon atoms;

R5 is independently R4 wherein each R4 is substituted with 0 to 3 R3 groups;

R5a is independently alkylene of 1 to 12 carbon atoms, alkenylene of 2 to 12 carbon
 atoms, or alkynylene of 2-12 carbon atoms any one of which alkylene, alkenylene or
 alkynylene is substituted with 0-3 R3 groups;

15 R6a is independently H or an ether- or ester-forming group;

R6b is independently H, a protecting group for amino or the residue of a carboxyl-
 containing compound;

R6c is independently H or the residue of an amino-containing compound;

W3 is W4 or W5;

20 W4 is R5, -C(Y1)R5, -C(Y1)W5, -SO2R5, or -SO2W5;

W5 is carbocycle or heterocycle wherein W5 is independently substituted with 0 to 3
 R2 groups;

m1 is independently an integer from 0 to 12, wherein the sum of all m1's within each
 individual claim of A1, A2 or A3 is 12 or less; and

25 m2 is independently an integer from 0 to 2.


149. The compound of claim 148 wherein:

A1 is -(C(R2)(R2))m1-W3, wherein W3 is substituted with 1 A3 group;

A2 is -(C(R2)(R2))m1-W3; and


30 A3 is -(C(R2)(R2))m1-P(Y1)(Y1R6a)(Y1R6a).

150. The compound of claim 148 having the formula:




5

151. The compound of claim 148 having the formula:




152. The compound of claim 151 having the formula:




10

153. The compound of claim 152 wherein W_3 is $-OC(O)NH_2$.

154. The compound of claim 152 having the formula:




155. The compound of claim 151 having the formula:



5


156. The compound of claim 155 wherein W_5 is a pyridine heterocycle bonded to $-C(R_2)_2-$ at the 2, 3, 4, 5 or 6 position.

157. The compound of claim 155 wherein W_5-A_3 has a formula selected from:




10

158. The compound of claim 155 wherein A_3 has a formula selected from:




wherein m_1 is 1, 2, 3, 4, 5, 6, 7 or 8, and the phenyl carbocycle is substituted with 0 to 3 R_2 groups.

- 5 159. The compound of claim 158 wherein A_3 has a formula selected from:



160. The compound of claim 159 wherein A_3 has a formula selected from:



10

161. A method of inhibiting the activity of HIV RT comprising the step of contacting

a sample suspected of containing HIV RT with a composition of claim 1.

162. The method of claim 161 wherein the HIV RT is *in vivo*.

5 163. A method for the treatment or prevention of the symptoms or effects of an HIV infection in an infected animal which comprises administering to said animal a formulation comprising a therapeutically effective amount of a compound according to claim 1.

10 164. The method of claim 163 wherein the compound is formulated with a pharmaceutically acceptable carrier.

15 165. A method of targeting white blood cells comprising administering to an HIV infected animal a formulation comprising a therapeutically effective amount of a compound according to claim 1 whereby the compound or metabolites derived from the compound accumulate in the white blood cells of the HIV infected animal.

166. The use of a compound of claim 1 to prepare a medicament for treatment of HIV infection.

20 167. The method of claim 163 wherein the formulation further comprises a second active ingredient selected from a protease inhibitor (Prt), a nucleoside reverse transcriptase inhibitor (NRTI), a non- nucleoside reverse transcriptase inhibitor (NNRTI), and an integrase inhibitor.

25 168. A process for preparing a compound of claim 1 wherein a compound comprising A³ or a precursor to A³ is reacted with an imidazole compound whereby a compound of claim 1 is formed.

30 169. A compound of the formula MBF.


170. In a non-nucleotide reverse transcriptase inhibitor, the improvement comprising a substituent having a phosphonate or phosphonate prodrug.

171. The improved non-nucleotide reverse transcriptase inhibitor compound of claim


5 170 selected from:

Capravirine-like phosphonate NNRTI compounds,
PETT-like phosphonate NNRTI compounds,
Pyrazole-like phosphonate NNRTI compounds,
Urea-PETT-like phosphonate NNRTI compounds,
10 Nevaripine-like phosphonate NNRTI compounds,
Quinazolinone-like phosphonate NNRTI compounds,
Efavirenz-like phosphonate NNRTI compounds,
Benzophenone-like phosphonate NNRTI compounds,
Pyrimidine-like phosphonate NNRTI compounds,
15 SJ3366-like phosphonate NNRTI compounds,
Delavirdine-like phosphonate NNRTI compounds,
Emivirine-like phosphonate NNRTI compounds,
Loviride-like phosphonate NNRTI compounds, and
UC781-like phosphonate NNRTI compounds;
20 and pharmaceutically acceptable salts, hydrates, and formulations thereof.





172. The improved non-nucleotide reverse transcriptase inhibitor compound of claim
170 of the Formulas:





25



where Z is CH or N,






where A is O, S, or N(R^x) and X is O or S;

5

wherein:


A⁰ is A¹, A² or W³ with the proviso that the compound includes at least one A¹;

A¹ is:




10

A² is:



A³ is:




Y^1 is independently O, S, $\text{N}(\text{R}^x)$, $\text{N}(\text{O})(\text{R}^x)$, $\text{N}(\text{OR}^x)$, $\text{N}(\text{O})(\text{OR}^x)$, or $\text{N}(\text{N}(\text{R}^x)(\text{R}^x))$;

Y^2 is independently a bond, O, $\text{N}(\text{R}^x)$, $\text{N}(\text{O})(\text{R}^x)$, $\text{N}(\text{OR}^x)$, $\text{N}(\text{O})(\text{OR}^x)$, $\text{N}(\text{N}(\text{R}^x)(\text{R}^x))$,

$-\text{S}(\text{O})_{\text{M2}}-$, or $-\text{S}(\text{O})_{\text{M2}}-\text{S}(\text{O})_{\text{M2}}-$;

5 R^x is independently H, R^1 , W^3 , a protecting group, or the formula:



R^y is independently H, W^3 , R^2 or a protecting group;

R^1 is independently H or an alkyl of 1 to 18 carbon atoms;

10 R^2 is independently H, R^1 , R^3 or R^4 wherein each R^4 is independently substituted with 0 to 3 R^3 groups, or taken together at a carbon atom, two R^2 groups form a ring of 3 to 8 carbons and the ring may be substituted with 0 to 3 R^3 groups;

R^3 is R^{3a} , R^{3b} , R^{3c} or R^{3d} , provided that when R^3 is bound to a heteroatom, then R^3 is R^{3c} or R^{3d} ;

R^{3a} is F, Cl, Br, I, $-\text{CN}$, N_3 or $-\text{NO}_2$;

15 R^{3b} is Y^1 ;

R^{3c} is $-\text{R}^x$, $-\text{N}(\text{R}^x)(\text{R}^x)$, $-\text{SR}^x$, $-\text{S}(\text{O})\text{R}^x$, $-\text{S}(\text{O})_2\text{R}^x$, $-\text{S}(\text{O})(\text{OR}^x)$, $-\text{S}(\text{O})_2(\text{OR}^x)$, $-\text{OC}(\text{Y}^1)\text{R}^x$, $-\text{OC}(\text{Y}^1)\text{OR}^x$, $-\text{OC}(\text{Y}^1)(\text{N}(\text{R}^x)(\text{R}^x))$, $-\text{SC}(\text{Y}^1)\text{R}^x$, $-\text{SC}(\text{Y}^1)\text{OR}^x$, $-\text{SC}(\text{Y}^1)(\text{N}(\text{R}^x)(\text{R}^x))$, $-\text{N}(\text{R}^x)\text{C}(\text{Y}^1)\text{R}^x$, $-\text{N}(\text{R}^x)\text{C}(\text{Y}^1)\text{OR}^x$, or $-\text{N}(\text{R}^x)\text{C}(\text{Y}^1)(\text{N}(\text{R}^x)(\text{R}^x))$;

R^{3d} is $-\text{C}(\text{Y}^1)\text{R}^x$, $-\text{C}(\text{Y}^1)\text{OR}^x$ or $-\text{C}(\text{Y}^1)(\text{N}(\text{R}^x)(\text{R}^x))$;

20 R^4 is an alkyl of 1 to 18 carbon atoms, alkenyl of 2 to 18 carbon atoms, or alkynyl of 2 to 18 carbon atoms;

R^5 is R^4 wherein each R^4 is substituted with 0 to 3 R^3 groups;

W^3 is W^4 or W^5 ;

W^4 is R^5 , $-\text{C}(\text{Y}^1)\text{R}^5$, $-\text{C}(\text{Y}^1)\text{W}^5$, $-\text{SO}_2\text{R}^5$, or $-\text{SO}_2\text{W}^5$;

W^5 is carbocycle or heterocycle wherein W^5 is independently substituted with 0 to 3 R^2 groups;

W^6 is W^3 independently substituted with 1, 2, or 3 A^3 groups;

W^7 is a heterocycle bonded through a nitrogen atom of said heterocycle and
 5 independently substituted with 0, 1 or 2 A^0 groups;

$M2$ is 0, 1 or 2;


$M12a$ is 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12;

$M12b$ is 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12;


$M1a$, $M1c$, and $M1d$ are independently 0 or 1; and

10 $M12c$ is 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12.


173. The improved non-nucleotide reverse transcriptase inhibitor compound of claim 172 of the Formulas:




Ia



Ib






Ic and




Id .

174. The improved non-nucleotide reverse transcriptase inhibitor compound of claim 172 of the Formulas:






and




5

175. The improved non-nucleotide reverse transcriptase inhibitor compound of claim 172 of the Formulas:




and




10


176. The improved non-nucleotide reverse transcriptase inhibitor compound of claim 172 of the Formulas:




5 177. The improved non-nucleotide reverse transcriptase inhibitor compound of claim 172 of the Formulas:




10 178. The improved non-nucleotide reverse transcriptase inhibitor compound of claim 172 of the Formulas:




179. The improved non-nucleotide reverse transcriptase inhibitor compound of claim 172 of the Formulas:




180. The improved non-nucleotide reverse transcriptase inhibitor compound of claim 172 of the Formulas:




181. The improved non-nucleotide reverse transcriptase inhibitor compound of claim 172 of the Formulas:




182. The improved non-nucleotide reverse transcriptase inhibitor compound of claim 172 of the Formulas:




5 183. The improved non-nucleotide reverse transcriptase inhibitor compound of claim 172 of the Formulas:



184. The improved non-nucleotide reverse transcriptase inhibitor compound of claim 172 of the Formulas:



5 185. The improved non-nucleotide reverse transcriptase inhibitor compound of claim 172 of the Formulas:




10 186. In a non-nucleotide reverse transcriptase inhibitor not containing a phosphonate or phosphonate prodrug, the improvement comprising a substituent having a phosphonate or phosphonate prodrug.






187. The improved non-nucleotide reverse transcriptase inhibitor compound of claim 186 selected from:

15 Capravirine-like phosphonate NNRTI compounds,
 PETT-like phosphonate NNRTI compounds,

Pyrazole-like phosphonate NNRTI compounds,
 Urea-PETT-like phosphonate NNRTI compounds,
 Nevaripine-like phosphonate NNRTI compounds,
 Quinazolinone-like phosphonate NNRTI compounds,
 5 Efavirenz-like phosphonate NNRTI compounds,
 Benzophenone-like phosphonate NNRTI compounds,
 Pyrimidine-like phosphonate NNRTI compounds,
 SJ3366-like phosphonate NNRTI compounds,
 Delavirdine-like phosphonate NNRTI compounds,
 10 Emivirine-like phosphonate NNRTI compounds,
 Loviride-like phosphonate NNRTI compounds, and
 UC781-like phosphonate NNRTI compounds;
 and pharmaceutically acceptable salts, hydrates, and formulations thereof.

15 188. The improved non-nucleotide reverse transcriptase inhibitor compound of claim
 186 of the Formulas:






where A is O, S, or N(R^x) and X is O or S;

wherein:


A^0 is A^1 , A^2 or W^3 with the proviso that the compound includes at least one A^1 ;

A^1 is:




5

A^2 is:




A^3 is:



- 10 Y^1 is independently O, S, $N(R^x)$, $N(O)(R^x)$, $N(OR^x)$, $N(O)(OR^x)$, or $N(N(R^x)(R^x))$;
 Y^2 is independently a bond, O, $N(R^x)$, $N(O)(R^x)$, $N(OR^x)$, $N(O)(OR^x)$, $N(N(R^x)(R^x))$,
 $-S(O)_{M2-}$, or $-S(O)_{M2-}S(O)_{M2-}$;

R^x is independently H, R^1 , W^3 , a protecting group, or the formula:



15

R^y is independently H, W^3 , R^2 or a protecting group;

R^1 is independently H or an alkyl of 1 to 18 carbon atoms;

R^2 is independently H, R^1 , R^3 or R^4 wherein each R^4 is independently substituted with 0 to 3 R^3 groups, or taken together at a carbon atom, two R^2 groups form a ring of 3 to 8 carbons and the ring may be substituted with 0 to 3 R^3 groups;

R^3 is R^{3a} , R^{3b} , R^{3c} or R^{3d} , provided that when R^3 is bound to a heteroatom, then R^3 is R^{3c} or R^{3d} ;

R^{3a} is F, Cl, Br, I, -CN, N_3 or $-NO_2$;

R^{3b} is Y^1 ;

R^{3c} is $-R^x$, $-N(R^x)(R^x)$, $-SR^x$, $-S(O)R^x$, $-S(O)_2R^x$, $-S(O)(OR^x)$, $-S(O)_2(OR^x)$, $-OC(Y^1)R^x$, $-OC(Y^1)OR^x$, $-OC(Y^1)(N(R^x)(R^x))$, $-SC(Y^1)R^x$, $-SC(Y^1)OR^x$, $-SC(Y^1)(N(R^x)(R^x))$, $-N(R^x)C(Y^1)R^x$, $-N(R^x)C(Y^1)OR^x$, or $-N(R^x)C(Y^1)(N(R^x)(R^x))$;

R^{3d} is $-C(Y^1)R^x$, $-C(Y^1)OR^x$ or $-C(Y^1)(N(R^x)(R^x))$;

R^4 is an alkyl of 1 to 18 carbon atoms, alkenyl of 2 to 18 carbon atoms, or alkynyl of 2 to 18 carbon atoms;

R^5 is R^4 wherein each R^4 is substituted with 0 to 3 R^3 groups;

W^3 is W^4 or W^5 ;

W^4 is R^5 , $-C(Y^1)R^5$, $-C(Y^1)W^5$, $-SO_2R^5$, or $-SO_2W^5$;

W^5 is carbocycle or heterocycle wherein W^5 is independently substituted with 0 to 3 R^2 groups;

W^6 is W^3 independently substituted with 1, 2, or 3 A^3 groups;

W^7 is a heterocycle bonded through a nitrogen atom of said heterocycle and independently substituted with 0, 1 or 2 A^0 groups;

$M2$ is 0, 1 or 2;




$M12a$ is 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12;

$M12b$ is 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12;

$M1a$, $M1c$, and $M1d$ are independently 0 or 1; and


$M12c$ is 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12.

189. The improved non-nucleotide reverse transcriptase inhibitor compound of claim 188 of the Formulas:




5

190. The improved non-nucleotide reverse transcriptase inhibitor compound of claim
10 188 of the Formulas:




191. The improved non-nucleotide reverse transcriptase inhibitor compound of claim 188 of the Formulas:


5




192. The improved non-nucleotide reverse transcriptase inhibitor compound of claim 188 of the Formulas:




193. The improved non-nucleotide reverse transcriptase inhibitor compound of claim 188 of the Formulas:




194. The improved non-nucleotide reverse transcriptase inhibitor compound of claim 188 of the Formulas:




195. The improved non-nucleotide reverse transcriptase inhibitor compound of claim 188 of the Formulas:




196. The improved non-nucleotide reverse transcriptase inhibitor compound of claim 188 of the Formulas:




197. The improved non-nucleotide reverse transcriptase inhibitor compound of claim 188 of the Formulas:




198. The improved non-nucleotide reverse transcriptase inhibitor compound of claim 188 of the Formulas:




5 199. The improved non-nucleotide reverse transcriptase inhibitor compound of claim 188 of the Formulas:



200. The improved non-nucleotide reverse transcriptase inhibitor compound of claim 188 of the Formulas:



5 201. The improved non-nucleotide reverse transcriptase inhibitor compound of claim 188 of the Formulas:



202. An MBF compound of Table 100.

203. A compound described herein.

204. A compound of claim 203 described in the schemes or examples.

205. A method of making a compound described herein.

206. A method of claim 205 described in the schemes or examples.

207. The use of a compound described here for treatment of HIV in humans.

5

208. The method of claim 207 wherein the compound is described in the schemes or examples.

209. The use of a compound described here in the manufacture of a medicament.

10


210. The use of claim 209 wherein the compound is described in the schemes or examples.

211. An non-nucleotide reverse transcriptase inhibitor compound capable of
15 accumulating in human PBMCs.

212. The compound of claim 211 further comprising a phosphonate or phosphonate prodrug.

20 213. The compound of claim 212 wherein the phosphonate or phosphonate prodrug
are of the formula A³:


A³ is:



Y¹ is independently O, S, N(R^x), N(O)(R^x), N(OR^x), N(O)(OR^x), or N(N(R^x)(R^x));

25 **Y² is independently a bond, O, N(R^x), N(O)(R^x), N(OR^x), N(O)(OR^x), N(N(R^x)(R^x)),
-S(O)_{M2}-, or -S(O)_{M2}-S(O)_{M2}-;**

R^x is independently H, R¹, W³, a protecting group, or the formula:



R^Y is independently H, W^3 , R^2 or a protecting group;

R^1 is independently H or an alkyl of 1 to 18 carbon atoms;

R^2 is independently H, R^1 , R^3 or R^4 wherein each R^4 is independently substituted with

5 0 to 3 R^3 groups;

R^3 is R^{3a} , R^{3b} , R^{3c} or R^{3d} , provided that when R^3 is bound to a heteroatom, then R^3 is R^{3c} or R^{3d} ;

R^{3a} is F, Cl, Br, I, -CN, N_3 or $-\text{NO}_2$;

R^{3b} is Y^1 ;

10 R^{3c} is $-\text{R}^x$, $-\text{N}(\text{R}^x)(\text{R}^x)$, $-\text{SR}^x$, $-\text{S}(\text{O})\text{R}^x$, $-\text{S}(\text{O})_2\text{R}^x$, $-\text{S}(\text{O})(\text{OR}^x)$, $-\text{S}(\text{O})_2(\text{OR}^x)$,

$-\text{OC}(\text{Y}^1)\text{R}^x$, $-\text{OC}(\text{Y}^1)\text{OR}^x$, $-\text{OC}(\text{Y}^1)(\text{N}(\text{R}^x)(\text{R}^x))$, $-\text{SC}(\text{Y}^1)\text{R}^x$, $-\text{SC}(\text{Y}^1)\text{OR}^x$,

$-\text{SC}(\text{Y}^1)(\text{N}(\text{R}^x)(\text{R}^x))$, $-\text{N}(\text{R}^x)\text{C}(\text{Y}^1)\text{R}^x$, $-\text{N}(\text{R}^x)\text{C}(\text{Y}^1)\text{OR}^x$, or $-\text{N}(\text{R}^x)\text{C}(\text{Y}^1)(\text{N}(\text{R}^x)(\text{R}^x))$;

R^{3d} is $-\text{C}(\text{Y}^1)\text{R}^x$, $-\text{C}(\text{Y}^1)\text{OR}^x$ or $-\text{C}(\text{Y}^1)(\text{N}(\text{R}^x)(\text{R}^x))$;

R^4 is an alkyl of 1 to 18 carbon atoms, alkenyl of 2 to 18 carbon atoms, or alkynyl of 2

15 to 18 carbon atoms;

R^5 is R^4 wherein each R^4 is substituted with 0 to 3 R^3 groups;

W^3 is W^4 or W^5 ;

W^4 is R^5 , $-\text{C}(\text{Y}^1)\text{R}^5$, $-\text{C}(\text{Y}^1)\text{W}^5$, $-\text{SO}_2\text{R}^5$, or $-\text{SO}_2\text{W}^5$;

W^5 is carbocycle or heterocycle wherein W^5 is independently substituted with 0 to 3 R^2

20 groups;

$\text{M}2$ is 0, 1 or 2;

$\text{M}12a$ is 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12;

$\text{M}12b$ is 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12;

$\text{M}1a$, $\text{M}1c$, and $\text{M}1d$ are independently 0 or 1; and

25 $\text{M}12c$ is 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12.

214. The compound of claim 213 wherein the intracellular half-life of the compound or an intracellular metabolite of the compound in human PBMCs is improved when compared to an analog of the compound not having the phosphonate or phosphonate prodrug.

5 215. The compound of claim 214 wherein the half-life is improved by at least about 50%.

216. The compound of claim 214 wherein the half-life is improved by at least about 100%.

10

217. The compound of claim 214 wherein the intracellular half-life of a metabolite of the compound in human PBMCs is improved when compared to an analog of the compound not having the phosphonate or phosphonate prodrug.

15 218. The compound of claim 217 wherein the half-life is improved by at least about 50%.

219. The compound of claim 217 wherein the half-life is improved by at least about 100%.

20

220. The compound of claim 217 wherein the half-life is improved by greater than 100%.

221. Use of a compound of the invention for the treatment of HIV infection.

25

222. Use of a compound of the invention in the manufacture of a medicament.

223. Use of a compound of the invention in the manufacture of a medicament for the treatment of disorders affecting white blood cells.

30

224. Method of treating a disorder affecting white blood cells, comprising:

administering a compound of the invention to a patient in need of white-blood-cell targeting.

225. Method of targeting a compound to white blood cells, comprising:

5 selecting a compound having a desired pharmaceutical activity and having a first structure;

modifying said first structure by replacing one or more atom of said first structure with an organic substituent comprising a phosphonate group or incipient phosphonate group to provide a compound having a second structure.

10 226. A method of manufacturing a non-nucleoside compound having both selectivity for white blood cells and a desired pharmaceutical activity, comprising:

chemically synthesizing a first molecule having a first structure containing a phosphonate or precursor phosphonate group, wherein said first structure differs from a
15 second structure of a compound known to have said desired pharmaceutical activity by having at least one hydrogen atom of said second structure replaced by an organic substituent comprising a phosphonate group or incipient phosphonate group.

20 227. The method of claim 226, wherein said first molecule is synthesized by a series of chemical reactions in which a hydrogen of said second structure is replaced by said organic substituent.

228. The method of claim 226, wherein said first molecule is synthesized by a series of chemical reactions that never includes a molecule of said second structure.

25 229. Method of accumulating an non-nucleotide reverse transcriptase inhibitor inside a white blood cell, comprising:

administering to a sample a composition comprising a compound of the invention.

30 230. The method of claim 229 wherein said sample is a patient.

231. The method of claim 229, wherein said compound of the invention has a

chemical structure A-B, wherein (a) a compound having structure A-H has non-nucleotide reverse transcriptase inhibitor activity and (b) substructure B comprises a phosphonate group or a precursor phosphonate group.

5 232. Method of increasing half-life of a non-nucleoside compound having anti-retroviral activity, comprising:

 replacing at least one hydrogen atom or organic radical of said compound by an organic substituent comprising a phosphonate group or incipient phosphonate.

10 233. Method of designing a drug having specificity for white blood cells for synthesis, comprising:

 obtaining a first list of first compounds having a desired activity;

 creating a second list of second compounds, each of said second compounds having a structure in which at least one hydrogen atom or organic radical of a compound of said first
15 list has been replaced by an organic substituent comprising a phosphonate group or incipient phosphonate group; and

 selecting a synthetic pathway capable of producing some or all of said second compounds from available starting materials, thereby providing a third list of compounds and associated synthetic techniques.

20

 234. Method of manufacturing a pharmaceutical composition having said specificity of claim 233, comprising:

 synthesizing a compound selected from said third list using said associated synthetic technique; and

25 admixing said synthesized compound with a pharmaceutically acceptable carrier.

 235. A composition produced by the method of claim 234.

 236. Method for producing a pharmaceutical composition having specificity for white
30 blood cells, comprising:

 admixing a therapeutically effective amount of a compound of the invention with a pharmaceutically acceptable carrier.